



My Truqap and fulvestrant treatment card

For my Truqap treatment, I take:

- ☐ Two 200 mg tablets
- ☐ Two 160 mg tablets
- ☐ One 200 mg tablet

At: (these times should be 12 hours apart)

..... am and pm

On: (please circle 4 consecutive days)

M : T : W : T : F : S : S

In my first month of fulvestrant treatment, I will have:
(please insert dates)

Injection 1:

Injection 2
(15 days later):

Injection 3
(29 days later):

DO NOT STOP OR CHANGE YOUR TREATMENT
UNLESS TOLD TO BY YOUR CANCER CARE TEAM.



If you have any questions or concerns about your treatment, speak to your cancer care team or scan this QR code to access the Patient Information Leaflet

Truqap may increase the risk of serious side effects, for example diabetic ketoacidosis (DKA). Immediately contact your doctor or seek urgent medical attention if you experience symptoms that may be related to DKA such as nausea, vomiting, stomach pain, trouble breathing, fruity breath, confusion, tiredness, or drowsiness.

Scan the QR code above for a complete list of side effects.

Reporting side effects

▼ This medicinal product is subject to additional monitoring.

This will allow quick identification of new safety information. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at <https://yellowcard.mhra.gov.uk/>.

GB-68306 | August 2025

This material was created and funded by AstraZeneca for patients taking this medicine.

**SHOW THIS CARD WHENEVER YOU SPEAK
TO A DOCTOR, NURSE OR PHARMACIST**